Further reading

Haaheim LR. Surveillance of the antibody status to current influenza strains in the Norwegian population. 1: Method of serum sampling and evaluation of the A/USSR/77 (H1N1) impact in various age groups in 1978. NIPH Ann. 1979 Dec;2(2):45-54. PMID: 548838.

Haaheim LR. Surveillance of the antibody status to current influenza strains in the Norwegian population. 2: Antibodies to the A/USSR/90/77 (H1N1) virus in various age groups after the second epidemic wave. NIPH Ann. 1980 May;3(1):49-56. PMID: 7454125.

Waalen K, Kilander A, Dudman SG, Ramos-Ocao R, Hungnes O.
Age-dependent prevalence of antibodies cross-reactive to the influenza A(H3N2) variant virus in sera collected in Norway in 2011.
Euro Surveill. 2012 May 10;17(19):20170. PMID: 22607964.

Fossum E, Rohringer A, Aune T, Rydland KM, Bragstad K, Hungnes O.
Antigenic drift and immunity gap explain reduction in protective responses against influenza A(H1N1)pdm09 and A(H3N2) viruses during the COVID-19 pandemic: a cross-sectional study of human sera collected in 2019, 2021, 2022, and 2023.
Virol J. 2024 Mar 6;21(1):57. doi: 10.1186/s12985-024-02326-w. Erratum in: Virol J. 2024 Mar 18;21(1):66. doi: 10.1186/s12985-024-02341-x. PMID: 38448981; PMCID: PMC10916265.

Waalen K, Kilander A, Dudman SG, Krogh GH, Aune T, Hungnes O.
High prevalence of antibodies to the 2009 pandemic influenza A(H1N1) virus in the Norwegian population following a major epidemic and a large vaccination campaign in autumn 2009.
Euro Surveill. 2010 Aug 5;15(31):19633. PMID: 20738992.

Norwegian reports to WHO influenza vaccine composition meetings:
https://www.fhi.no/ss/influensa/sesonginfluensa/who-rapporter/